



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Statistical reports of States and cities of the United States—Yearly and monthly.

COLORADO—*Denver*.—Month of March, 1902. Estimated population, 150,000. Total number of deaths, 215, including diphtheria, 1; enteric fever, 3; scarlet fever, 4, and 52 from tuberculosis.

Month of April, 1902. Total number of deaths, 202, including diphtheria, 4; enteric fever, 3; scarlet fever, 5; whooping cough, 3, and 51 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended May 24, 1902, from 91 observers, indicate that inflammation of bowels, erysipelas, and meningitis were more prevalent and pneumonia, intermittent fever, whooping cough, remittent fever, and diphtheria were less prevalent than in the preceding week. Meningitis was reported present at 9, whooping cough at 17, diphtheria at 22, enteric fever at 38, measles at 61, scarlet fever at 74, smallpox at 87, and phthisis pulmonalis at 223 places.

MISSOURI—*St. Joseph*.—Month of April, 1902. Estimated population, 105,000. Total number of deaths, 77, including diphtheria, 1; measles, 15; whooping cough, 1; smallpox, 2, and 8 from tuberculosis.

St. Louis.—Month of April, 1902. Estimated population, 598,000—white, 560,000; colored, 38,000. Total number of deaths, 991—white, 851; colored, 140—including diphtheria, 7; enteric fever, 14; scarlet fever, 17; whooping cough, 20; smallpox, 4, and 150 from phthisis pulmonalis.

NEW JERSEY—*Passaic*.—Two weeks ended May 24, 1902. Census population, 27,777. Total number of deaths, 26, including scarlet fever, 1, and 4 from measles.

OHIO—*Cleveland*.—Month of April, 1902. Estimated population, 390,000. Total number of deaths, 513, including diphtheria, 11; enteric fever, 6; measles, 3; whooping cough, 3; smallpox, 1, and 43 from tuberculosis.

East Liverpool.—Month of April, 1902. Census population, 16,486. Total number of deaths, 25, including measles, 1, and 2 from enteric fever.

NEW YORK—*Middletown*.—Month of April, 1902. Estimated population, 15,000. Total number of deaths, 21. No deaths from contagious diseases.

Report of immigration at Baltimore for the week ended May 31, 1902.

OFFICE OF U. S. COMMISSIONER OF IMMIGRATION,

Port of Baltimore, Md., May 31, 1902.

Number of alien immigrants who arrived at this port during the week ended May 31, 1902; also, names of vessels and ports from which they came. May 26. steamship *Barnstable*, from Port Antonio, with 7 immigrants; May 29. steamship *Dresden*, from Bremen, with 1,526 immigrants. Total, 1,533.

BERTRAM M. STUMP,
Acting Commissioner.